

	<b>Evidence</b>	the actual evidence for it. For example, if you feel that you never do anything right, you could list several things you have done well recently.
1.	<b>The Double-Standard Method</b>	Instead of putting yourself down in a harsh, condemning way, talk to yourself in the same compassionate way you would talk to a friend with a similar problem.
4.	<b>The Experimental Technique</b>	Do an experiment to test the validity of your negative thought. For example, if during the episode of panic, you become terrified that you're about to die of a heart attack, you could jog or run up and down several flights of stairs. This will prove that your heart is healthy and strong.
1.	<b>Thinking in Shades of Grey</b>	Although this method might sound bleak, the effects can be illuminating. Instead of thinking about your problems in almost nothing extremes, evaluate things on a range of 0 to 100. When things don't work out as well as you hoped, think about the experience as a partial success rather than a complete failure. See what you can learn from the situation.
4.	<b>The Survey Method</b>	Ask people questions to find out if your thoughts and attitudes are realistic. For example, if you believe that public speaking anxiety is abnormal and shameful, ask several friends if they ever felt nervous before they gave a talk.
1.	<b>Define Terms</b>	When you label yourself "inferior" or "a fool" or "a loser," ask, "What is the definition of a 'fool'?" You will feel better when you see that there is no such thing as a "fool" or a "loser."
4.	<b>The Semantic Method</b>	Simply substitute language that is less colorful and emotionally loaded. This method is helpful for "should statements." Instead of telling yourself "I shouldn't have made that mistake," you can say, "It would be better if I hadn't made that mistake."
4.	<b>No attribution</b>	Instead of automatically assuming that you are "bad" and blaming yourself entirely for a problem, think about the many factors that