Subtracting Mixed Fractions and Borrowing (Different Denominator Work)

A) When the denominators of the fraction you are subtracting are different, you must find the lowest common denominator. Using the example, you must find the lowest common denominator of 5 and 10. The lowest common denominator or least common multiple os 10. We find the LCM by counting the multiples of both denominators.

5 - 5, 10, 15, 20
The LCM of 5 and 10 is 10!

B) After you find your LCM, rewrite both fractions. Because one of the had the LCM for a denominator, you simply copy that fraction over. The denominator of five has to be turned into the LCM. Because you have to multiply the denominator of 5 by 2 to get 10, you also have to increase the denominator by the same factor. Two fifths turns into four tenths.

d) The top fraction is now large enough to be subtracted from!

1)
$$16\frac{1}{2}$$
 $- 5\frac{4}{7}$

2)
$$12\frac{1}{6}$$
 $- 9\frac{3}{4}$

3)
$$14\frac{4}{9}$$
 - $11\frac{7}{18}$

4)
$$4 \frac{4}{5}$$
 $- 3 \frac{2}{10}$

5)
$$19 \frac{1}{6}$$
 $- 6 \frac{4}{9}$

6)
$$32 \frac{1}{3}$$
 $-24 \frac{1}{5}$

7)
$$24 \frac{3}{16}$$
 $- 16 \frac{1}{4}$

8)
$$9 \frac{3}{6} - 6 \frac{2}{3}$$

9)
$$64 \frac{1}{3}$$
 $-45 \frac{4}{9}$

10)
$$32\frac{1}{6}$$

$$- 10\frac{2}{12}$$

9)
$$64\frac{1}{3}$$
 10) $32\frac{1}{6}$ 11) $86\frac{3}{8}$ $-45\frac{4}{9}$ $-10\frac{2}{12}$ $-83\frac{5}{16}$

12)
$$4 \frac{1}{6}$$
 $- 2 \frac{5}{24}$