Conservation of Energy and Momentum Worksheet

Recall the basic relations:

$$\begin{split} E_i &= E_f \\ P_i &= P_f \\ \Delta E \Delta t \geq \frac{h}{2\pi} \\ \Delta p \Delta x \geq \frac{h}{2\pi} \end{split}$$

Equations:

1.
$$E = K + U + mc^2$$
 (usually $U = 0$)

3.
$$E = hf$$
 (for a photon)

4.
$$p = mv$$
 (for a massive particle at low speeds

5.
$$p = \frac{E}{c}$$
 (for a massless sparticle)

Constants:

$$h = 6.63 \times 10^{-34} \,\text{J} \cdot \text{s} = 4.14 \times 10^{-21} \,\text{MeV} \cdot \text{s}$$

$$c = 3.00 \times 10^8 \, \text{m/s}$$

Mass of electron = mass of positron =
$$9.1 \times 10^{-31}$$
 kg = $.511$ MeV/c²

Mass of hydrogen atom =
$$1.67 \times 10^{-27}$$
 kg

Example 1:

A gamma ray photon has energy equal to 5 MeV. What is its:
a. momentum (in kg-m/s)?
b. momentum (in MeV/c)?

- c. frequency (in s⁻¹)?

Example 2:

An electron and positron are at rest in a particular frame of reference. What is their:

- a. total energy (in J)?
- b. total energy (in MeV)?

Example 3 A 5 MeV gamma ray turns into an electron-positron pair moving in the same direction as the gamma ray. What is:

- a. the total momentum of the electron and positron (in MeV/c)?
 b. the total energy of the electron and positron (in MeV) using energy conservation?
- c. the total energy of the electron and positron using Equation 2 (above) (in MeV)?
 d. the difference in energy between the results of (b) and (c)?

Example 4:

Using the result (d) from Example 3, and the uncertainty principle, for how much time can the electron-positron pair exist before they must annihilate (turn back into a gamma ray)?