

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are

■ Find the values of the unknowns.
for $\sin A$, $\cos A$,
sec A , $\csc A$,
tan A , and cot A .
Angle A is in standard position.
Coordinates of P are $(-3, 4)$.

Length of r is
Coordinates of P are
Angle A is
Coordinates of P are