


# TRIG CHEAT SHEET

## DEFINITION OF THE TRIG FUNCTIONS

### Right Triangle Definition

For this definition we assume that


$$0 < \theta < \frac{\pi}{2} \text{ or } 0^\circ < \theta < 90^\circ.$$



$$\begin{aligned} \sin \theta &= \frac{\text{opposite}}{\text{hypotenuse}} & \csc \theta &= \frac{\text{hypotenuse}}{\text{opposite}} \\ \cos \theta &= \frac{\text{adjacent}}{\text{hypotenuse}} & \sec \theta &= \frac{\text{hypotenuse}}{\text{adjacent}} \\ \tan \theta &= \frac{\text{opposite}}{\text{adjacent}} & \cot \theta &= \frac{\text{adjacent}}{\text{opposite}} \end{aligned}$$

### Unit Circle Definition

For this definition  $\theta$  is any angle.



$$\begin{aligned} \sin \theta &= \frac{y}{1} = y & \csc \theta &= \frac{1}{y} \\ \cos \theta &= \frac{x}{1} = x & \sec \theta &= \frac{1}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y} \end{aligned}$$

## FACTS AND PROPERTIES

### Domain

The domain is all the values of  $\theta$  that can be plugged into the function.

$$\begin{aligned} \sin \theta, \quad \theta &\text{ can be any angle} \\ \cos \theta, \quad \theta &\text{ can be any angle} \\ \tan \theta, \quad \theta &\neq \left(n + \frac{1}{2}\right)\pi, \quad n = 0, \pm 1, \pm 2, \dots \\ \csc \theta, \quad \theta &\neq n\pi, \quad n = 0, \pm 1, \pm 2, \dots \\ \sec \theta, \quad \theta &\neq \left(n + \frac{1}{2}\right)\pi, \quad n = 0, \pm 1, \pm 2, \dots \\ \cot \theta, \quad \theta &\neq n\pi, \quad n = 0, \pm 1, \pm 2, \dots \end{aligned}$$

### Range

The range is all possible values to get out of the function.

$$\begin{aligned} -1 \leq \sin \theta \leq 1 & & \csc \theta \geq 1 \text{ and } \csc \theta \leq -1 \\ -1 \leq \cos \theta \leq 1 & & \sec \theta \geq 1 \text{ and } \sec \theta \leq -1 \\ -\infty < \tan \theta < \infty & & -\infty < \cot \theta < \infty \end{aligned}$$

### Period

The period of a function is the number,  $T$ , such that  $f(\theta + T) = f(\theta)$ . So, if  $\omega$  is a fixed number and  $\theta$  is any angle we have the following periods.

$$\begin{aligned} \sin(\omega\theta) &\rightarrow T = \frac{2\pi}{\omega} \\ \cos(\omega\theta) &\rightarrow T = \frac{2\pi}{\omega} \\ \tan(\omega\theta) &\rightarrow T = \frac{\pi}{\omega} \\ \csc(\omega\theta) &\rightarrow T = \frac{2\pi}{\omega} \\ \sec(\omega\theta) &\rightarrow T = \frac{2\pi}{\omega} \\ \cot(\omega\theta) &\rightarrow T = \frac{\pi}{\omega} \end{aligned}$$