

LARSON ALGEBRA 2
CHAPTER 1, LESSON 3, EXTRA EXAMPLES

Extra Example 1 Solving an Equation with a Variable on One Side

Solve $\frac{2}{9}x + 8 = 16$.

SOLUTION

Your goal is to isolate the variable on one side of the equation.

$$\frac{2}{9}x + 8 = 16$$

Write original equation.

$$\frac{2}{9}x = 8$$

Subtract 8 from each side.

$$x = \frac{9}{2}(8)$$

Multiply each side by $\frac{9}{2}$, the reciprocal of $\frac{2}{9}$.

$$x = 36$$

Simplify.

◆ The solution is 36.

Check : Check $x = 36$ in the original equation.

$$\frac{2}{9}(36) + 8 = 16$$

Substitute **36** for x .

$$16 = 16 \checkmark$$

Solution checks.

Extra Example 2 Solving an Equation with a Variable on Both Sides

Solve $12n - 3 = 4n + 21$.

SOLUTION

$$12n - 3 = 4n + 21$$

Write original equation.

$$8n - 3 = 21$$

Subtract $4n$ from each side.

$$8n = 24$$

Add 3 to each side.

$$n = 3$$

Divide each side by 8.

◆ The solution is 3. Check this in the original equation.