MEDICATION SHEET (Pre-Op) Allergies: Codeine

Generic/Trade Classification	Route/ Dose	Freq./ Time	Action	Purpose, Nursing Implications Major Side Effects
Cefazolin Ancef Anti-infectives	IV 1g	30 to 60 minutes before surgery to, q 8 hrs up to 24 hrs.	Perioperative prophylaxis	Avoid infections, Monitor IV site freq. (pain, redness, swelling) Observe pt. for S/S of rash, pruritus, laryngeal edema or wheezing. D/C med and call Dr.
Sublimaze Fentanyl Opioid Analgesics	IV 1-2 mg	45 minutes Before surgery	Binds to opiate receptors the CNS, altering the response to the perception of pain.	Supplements in anesthesia – dilute in 0.9% NaCl Closely Monitor RR, B/P Caution the pt. change position slowly Educate pt. no driving for 24 hrs. and avoid alcohol. Reversal (Narcan)
Metoclopramide Reglan Antiemetics	IV 10mg	Per/post operative	Treatment & prevention of nausea and vomiting	Decreases nausea & vomiting Assess pt. of S/S N/V, Assess bowel sounds before and after. Administer slow to avoid side effects such as: anxiety & restlessness.
Glycoryrolate Robinul Anticholinergics	IV 4.4mch/kg	30-60 minutes preop	Decrease Gi & Respiratory secretions	Reduces secretions Decreases risk for aspiration, decreases risk vomiting Used in larger case

Ht.: <u>5' 3"</u> Wt.: <u>145</u>

Parenteral Infusions: 1000 ml of 0.9% NaCl